




Water is an increasingly scarce resource and is essential for Arizona's future. With Arizona's population growth and continued drought, citizens and water managers have been taking a closer look at water supplies in the state. Municipal, industrial, and agricultural water users are well-represented demand sectors, but water supplies and management to benefit the environment are not often considered. This bulletin explains the water demands of the environment in the Central Arizona Region, an area that includes the Verde River, Agua Fria and Upper Hassayampa groundwater basins, as well as the Prescott Phoenix, and Pinal Active Management Areas (AMAs).




Data Source: USGS stream gage data

**Figure 1. Elements of Environmental Flow Occurring in Seasonal Hydrographs**

This Central Arizona Region bulletin also introduces information essential for considering environmental water demands in discussions about water management. Environmental water demands (or environmental flow) refers to how much water a freshwater ecosystem needs to sustain itself. Arizona's native animals and plants are dependent on dynamic flows, which are commonly described according to five elements: magnitude, duration, frequency, timing and rate of change. For example, seasonal flood events (e.g. timing) and constant flows (e.g. duration) cue important biological events, like reproduction. The five elements of environmental flows are displayed in Figure 1 through a hydrograph of the San Pedro River's flows over the course of a year.

To consider the environment alongside other water sectors, we must first study the water demands of ecosystems. In Figure 2 the streams where we have quantified the current amount of streamflow that supports the environment (gray lines) and environmental water demands (black lines) are displayed in relation to key surface water resources. This region contains perennial (those that flow year-round) and intermittent (those that flow only part of the year) streams, riparian areas, and many major springs.




**Figure 2. Streams with Quantified Flows/Demands and Surface Water Resources in the Central Arizona Region**

Humans have an interconnected and dependent relationship with the environment. Nature provides recreation opportunities, economic benefits, and water supplies to sustain our communities. For example, an estimated \$150 million per year of the economy in the Verde Valley depends on water resources (Limbrunner et al, 2011).

How water is used in the Central Arizona Region is shown in Figure 3 by comparing the relative scale of human water demands by sector to existing minimum, median, and maximum flows available in the environment. The total size of the pie chart of human demands (far right) reflects the 3.3 million acre-feet annually withdrawn by all sectors (municipal, industrial, and agricultural) by source in the region. Median annual flows for the gaged rivers in the region are about one third the amount used by all human demand sectors, while maximum flood flows are two times greater. Surface water imported from the Colorado River is included in human surface water use but not in the current stream-flow quantities shown on Figure 3. Although human and environmental demands are not always mutually exclusive, some streams in the region no longer contain perennial flows because of human water uses.

**Figure 3: Human Demand and Current Flow in the Central Arizona Region (circle size indicates relative amount of water)**




\*In 2006 an additional 0.19 maf of effluent was also used to meet demand

Data Sources: ADWR 2010 (streamflow as measured by stream flow gages), WRDC 2011 (human demand)

Figure 4 shows the Central Arizona Region’s median streamflow as a single “stream” and how it interacts with groundwater and human demands. Flows into the environment are represented by blue arrows, while outflows to human and environmental demands are marked by green arrows. Note that all human sectors return some water to the environment after use. Also, water traveling through a river to farming or domestic uses downstream can support aquatic and riparian ecosystems (streamside) along the way. These connections between environmental and human demands can create opportunities for water management that is mutually beneficial.

**Figure 4: Water Demand and Use in the Central Arizona Region (Arrows indicate relative size of demand and recharge)**



Data Source: WRDC 2011

In the Central Arizona Region more stream reaches have intermittent flow (77%) than perennial flow (23%), (see Table 1a). Perennial streams are better understood than intermittent streams, but only half of all perennial streams have been studied. There are 18 known studies (16 quantitative and two qualitative) in this region that characterize some aspect of environmental water demands. These studies provide information on the flow velocity, water quality, and depth to groundwater needed to support native species.

Table 1b categorizes available information for select streams in the region by the elements of flow that have been studied. Although some Central Arizona streams have been studied for five flow elements, these studies focus on the demands of a few riparian species and do not address the flow demands and responses for the whole ecosystem. Fourteen of the 16 quantitative studies in this region examined multiple species' needs, and six of them quantified both environmental flow demands and ecological responses to flow. Only one study, on

**Table 1a: Data Coverage for Flow Demands in Central Region Streams**


% Area Perennial/ Intermittent	% Total Studied	% Perennial Streams Studied	% Intermittent Streams Studied
<i>(Miles)</i>			
23%/77%	16%	50%	5%
(600/1900)	(400)	(300)	(100)

**Table 1b: Flow Components Studied and Information Gaps for Perennial Streams in the Central Region**

River Name	Magnitude	Duration	Frequency	Timing of Flow	Rate of Change	Water Quality*
	<i>(% of the Stream Reach Studied)</i>					
Agua Fria River	S (22%)	NS	NS	S (22%)	NS	NS
E. Verde River	S	NS	NS	S	NS	NS
Gila River	S (6%)	S (6%)	S (6%)	S (6%)	S (6%)	S (6%)
Hassayampa River	S	S	S	S	S	S
Oak Creek	S	NS	S	S	NS	NS
Salt River	S	NS	NS	NS	NS	S
Santa Cruz River	S	NS	NS	NS	NS	NS
Sycamore Creek	S	S	S	S	NS	NS
Verde River	S	S	S	S	S	S

\* Does not include studies of water quality alone, these studies were not reviewed for this report. **S** = Entire stream surveyed, **S** = Reach (% of stream surveyed), **NS** = Not surveyed

Data Source: Nadeau and Megdal 2011



**Figure 3: Environmental Resources and Designations in the Central Arizona Region**

the East Verde River, fully describes the flows needed to maintain riparian ecosystem function in a local stream. Records from 2011 and early 2012 indicate that flows on the East Verde fluctuate around the recommended minimum flows, sometimes dipping below the level of flow needed to maintain the ecosystem.

Official designations by the state and/or federal government are made to protect stream reaches with high environmental values. These designations include Wild and Scenic Rivers, Instream Flow Permits and Applications, Arizona Department of Environmental Quality Unique Waters, Endangered Species Act Critical Habitat, and Federal Conservation Lands such as National Forest Wilderness or National Parks. Many stream reaches, such as on the Verde River, have multiple designations (see Figure 3 for the number of designations on stream reaches in this region). Having many designations on one reach can be an indication of an area with significant environmental resources. Different designations provide different types of protections for environmental flows, but having three designations does not necessarily mean the reach is better protected than a reach with one designation.



Statewide, ecosystem-level flow requirements remain poorly understood. Small scale studies that prescribe flows for a single reach exist for some areas, but cannot be applied across basins or regions. Two areas of agreement have emerged from studies done across the state: (1) riparian areas need both access to sufficient groundwater and carefully-timed flood flows to maintain water levels for established plants and for new plant growth; and (2) change to any element of flow can impact Arizona's aquatic and riparian ecosystems if flows are altered beyond the range of tolerance of native species.



*Verde River, Arizona. Photo Credit: Arizona State Parks*

The Central Arizona Region has a wealth of natural resources in its streams, springs, and riparian areas. Water demands of the environment in this region have been studied more frequently than in other areas of the state, but are still incomplete. Even with the available information, this region lacks clear, measurable management objectives for all of its streams.

Information available in the region on the relationships between components of flow and biological factors can be used for considering potential impacts of future water decisions. By comparing various environmental flow demands, such as species-specific water demands, with current conditions, areas needing protection or restoration can be identified.

These pages present a brief overview of the information available for the Central Arizona Region; more detailed information to help inform planning efforts throughout this region is available by contacting the WRRC.

## How you can apply this information

Those working to address the demands of all water sectors in Arizona can apply this information to:

- Determine how environmental flows interact with other demand sectors regionally,
- Identify factors putting environmental flow demands at risk,
- Identify studies needed to address key information gaps about environmental flows,
- Determine local priorities for ecosystems, and then identify water needed to preserve or restore those,
- Develop scenario analyses for water planning that incorporate the environment, and
- Share the information widely to increase understanding of regional resources and challenges.

## Contact Info

For assistance applying information about environmental water uses and needs in water planning, questions about methods used to create this bulletin or requests for our environmental water demand data please contact:

**Kelly Mott Lacroix**

**Email: [klacroix@email.arizona.edu](mailto:klacroix@email.arizona.edu)**

**Phone: (520) 621-3826**

The WRRC offers public presentations about this information as well as direct support for water planning processes as part of our Connecting the Environment to Arizona Water Planning (EnWaP) project.

[wrrc.arizona.edu](http://wrrc.arizona.edu)



## References

- Arizona Department of Water Resources (ADWR). (2010) "Arizona Water Atlas: Volume 1 Executive Summary." Phoenix, Arizona: ADWR, Office of Water Management.
- \_\_\_\_\_. (2009) "Arizona Major and Minor Springs." GIS Data File.
- Arizona Game and Fish Department (AGFD). (1997) Intermittent Streams GIS Data File.
- \_\_\_\_\_. (1993) "Riparian Vegetation." GIS Data File.
- Limbrunner, J., Sheer, D., Heberger, M., Cohen, M., Henderson, J., Raucher, B. (2011) "Policy Options for Water Management in the Verde Valley, Arizona." The Nature Conservancy, Flagstaff, AZ.
- Lowry, J. H. Jr., Ramsey, R. D., Thomas, K. A., Schrupp, D., Kepner, W., Sajwaj, T., Thompson, B. (2007) "Land cover classification and mapping." In J.S. Prior- Magee, et al., (Eds.) Southwest Regional Gap Analysis Final Report (Chapter 2). Moscow, ID: U.S. Geological Survey, Gap Analysis Program.
- Nadeau, J; S.B. Megdal. (2011) "Arizona Environmental Water Needs Assessment Report and Methodology Guidebook." Water Resources Research Center.
- The Nature Conservancy. (2010) "Arizona Freshwater Assessment: GIS Package."
- Water Resources Development Commission (WRDC). (2011) "Environmental Working Group - Arizona's Inventory of Water-Dependent Natural Resources."

*Bulletin prepared by: Brittany Choate, Leah Edwards, Kelly Mott Lacroix and Joanna B. Nadeau*